

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 136 493 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
26.09.2001 Bulletin 2001/39

(21) Application number: 00400803.3

(22) Date of filing: 23.03.2000

(51) Int Cl.7: C07D 495/04, A61K 31/513,
C07D 471/04, C07D 401/14,
C07D 491/04, A61P 25/28,
A61P 25/16
// (C07D495/04, 333:00,
221:00),
(C07D471/04, 221:00, 209:00),
(C07D491/04, 307:00, 221:00)

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(71) Applicants:
• SANOFI-SYNTHELABO
75013 Paris (FR)
• Mitsubishi-Tokyo Pharmaceuticals, Inc.
Chuo-ku, Tokyo 103-8405 (JP)


(72) Inventors:
• Almarlo-Garcia, Antonio
92290 Chatenay-Malabry (FR)
• Frost, Jonathan Reid
91320 Wissous (FR)
• Li, Adrien-Tak
92260 Fontenay-aux-Roses (FR)

(74) Representative: Thouret-Lemaître, Elisabeth et al
Sanofi-Synthelabo,
Département Brevets,
174, avenue de France
75013 Paris (FR)

(54) 2-(Thienopyridinyl)pyrimidone, 2-(furopyridinyl)pyrimidone 2-(isoquinolinyl)pyrimidone,
2-(pyridoindolyl)pyrimidone and 2-(benzofuropyridinyl)pyrimidone derivatives


(57) A pyrimidone derivative represented by formula
(I) or a salt thereof:

and The heterocyclic ring of formula :



wherein:

R1 represents a pyridyl ring optionally substituted by a
C₁₋₄ alkyl group, C₁₋₄ alkoxy group or a halogen atom,



is selected from the group : 2,3,4,9-tetrahydro-1H-pyri-
do[3,4-b]indole, 1,2,3,4-tetrahydroisoquinoline,
4,5,6,7-tetrahydrothieno[2,3-c]pyridine or 4,5,6,7-tet-
rahydrofuro[2,3-c]pyridine.

And a medicament comprising the said derivative
or a salt thereof as an active ingredient which is used
for preventive and/or therapeutic treatment of a neuro-
degenerative disease caused by abnormal activity of
GSK3 β such as Alzheimer's disease, Parkinson's dis-
ease, frontoparietal dementia, corticobasal degenera-
tion, Pick's disease, cerebrovascular accidents, brain
and spinal trauma, and peripheral neuropathies.

Description

Technical Field

- 5 [0001] The present invention relates to compounds that are useful as an active ingredient of a medicament for preventive and/or therapeutic treatment of neurodegenerative diseases caused by abnormal activity of GSK3.

Background Art

- 10 [0002] GSK3 β (glycogen synthase kinase 3 β) is a proline directed serine, threonine kinase that plays an important role in the control of metabolism, differentiation and survival. It was initially identified as an enzyme able to phosphorylate and hence inhibit glycogen synthase. It was later recognized that GSK3 β was identical to tau protein kinase 1 (TPK1), an enzyme that phosphorylates tau protein in epitopes that are also found to be hyperphosphorylated in Alzheimer's disease and in several tauopathies.

- 15 Interestingly, protein kinase B (AKT) phosphorylation of GSK3 β results in a loss of its kinase activity, and it has been hypothesized that this inhibition may mediate some of the effects of neurotrophic factors. Moreover, phosphorylation by GSK3 β of β -catenin, a protein involved in cell survival, results in its degradation by an ubiquitination dependent proteasome pathway.


- 20 Thus, it appears that inhibition of GSK3 β activity may result in neurotrophic activity. Indeed there is evidence that lithium, an uncompetitive inhibitor of GSK3 β , enhances neuriteogenesis in some models and also increases neuronal survival, through the induction of survival factors such as Bcl-2 and the inhibition of the expression of proapoptotic factors such as P53 and Bax.

- 25 Recent studies have demonstrated that β -amyloid increases the GSK3 β activity and tau protein phosphorylation. Moreover, this hyperphosphorylation as well as the neurotoxic effects of β -amyloid are blocked by lithium chloride and by a GSK3 β antisense mRNA. These observations strongly suggest that GSK3 β may be the link between the two major pathological processes in Alzheimer's disease : abnormal APP (Amyloid Precursor Protein) processing and tau protein hyperphosphorylation.

- 30 [0003] Although tau hyperphosphorylation results in a destabilization of the neuronal cytoskeleton, the pathological consequences of abnormal GSK3 β activity are, most likely, not only due to a pathological phosphorylation of tau protein because, as mentioned above, an excessive activity of this kinase may affect survival through the modulation of the expression of apoptotic and antiapoptotic factors. Moreover, it has been shown that β -amyloid-induced increase in GSK3 β activity results in the phosphorylation and, hence the inhibition of pyruvate dehydrogenase, a pivotal enzyme in energy production and acetylcholine synthesis.

- 35 [0004] Altogether these experimental observations indicate that GSK3 β may find application in the treatment of the neuropathological consequences and the cognitive and attention deficits associated to Alzheimer's disease, as well as to other acute and chronic neurodegenerative diseases. These include, in a nonlimiting manner, Parkinson's disease, frontoparietal dementia, corticobasal degeneration, Pick's disease, cerebrovascular accidents, peripheral neuropathies and brain and spinal cord trauma.

- 40 [0005] PCT application WO 98/24782 discloses compounds represented by the following formula (A):



55 wherein R represents a 2,6-dichlorobenzyl group, a 2-(2-chlorophenyl)ethylamino group, a 3-phenylpropylamino group, or a 1-methyl-3-phenylpropylamino group. The compounds represented by formula (A) are characterized by a 4-fluorophenyl group at the 5-position of the pyrimidine ring. The main pharmacological activity disclosed for the compounds


represented by formula (A) is an anti-inflammatory effect, whereas the compounds of the present invention represented by formula (I) herein below are useful as GSK3 β inhibitors or as medicaments for the therapeutic treatment of neurodegenerative diseases, and therefore, their pharmacological activities are totally different.

Disclosure of the Invention

[0006] An object of the present invention is to provide compounds useful as an active ingredient of a medicament for preventive and/or therapeutic treatment of neurodegenerative diseases. More specifically, the object is to provide novel compounds useful as an active ingredient of a medicament that enables prevention and/or treatment of the neurodegenerative diseases such as Alzheimer's disease.


Thus, the inventors of the present invention have identified compounds possessing inhibitory activity against GSK3 β . As a result, they found that compounds represented by the following formula (I) had the desired activity and were useful as an active ingredient of a medicament for preventive and/or therapeutic treatment of the aforementioned diseases.

[0007] The present invention thus provides pyrimidone derivatives represented by formula (I) or salts thereof, solvates thereof or hydrates thereof:




wherein:

- * R1 represents a pyridyl ring optionally substituted by a C₁₋₄ alkyl group, C₁₋₄ alkoxy group or a halogen atom ;
- * The heterocyclic ring of formula :



is selected from the group consisting of:



the heterocyclic rings being optionally substituted by 1 to 3 substituents selected from a C₁₋₆ alkyl group, halogeno, a C₁₋₂ perhalogenated alkyl group, a C₁₋₃ halogenated alkyl group, a hydroxyl group, a C₁₋₆ alkoxy group, methylenedioxy group, a nitro, a cyano, or a phenyl group; and

- * R₂ represents a hydrogen atom or a C₁₋₆ alkyl group.

[0008] According to another aspect of the present invention, there is provided a medicament comprising as an active ingredient a substance selected from the group consisting of the pyrimidone derivatives represented by formula (I) and the physiologically acceptable salts thereof, and the solvates thereof and the hydrates thereof. As preferred embodiments of the medicament, there are provided the aforementioned medicament which is used for preventive and/or therapeutic treatment of diseases caused by abnormal GSK3 β activity, and the aforementioned medicament which is used for preventive and/or therapeutic treatment of neurodegenerative diseases. As further preferred embodiments of the present invention, there are provided the aforementioned medicament wherein the diseases are selected from the group consisting of Alzheimer's disease, Parkinson's disease, frontoparietal dementia, corticobasal degeneration, Pick's disease, cerebrovascular accidents, brain and spinal cord trauma, and peripheral neuropathies; and the aforementioned medicament in the form of pharmaceutical composition containing the above substance as an active ingredient together with one or more pharmaceutical additives.

[0009] The present invention further provides an inhibitor of GSK3 β activity comprising as an active ingredient a substance selected from the group consisting of the pyrimidone derivatives of formula (I) and the salts thereof, and the solvates thereof and the hydrates thereof.

[0010] According to further aspects of the present invention, there are provided a method for preventive and/or therapeutic treatment of neurodegenerative diseases caused by abnormal GSK3 β activity, which comprises the step of administering to a patient a preventively and/or therapeutically effective amount of a substance selected from the group consisting of the pyrimidone derivatives of formula (I) and the physiologically acceptable salts thereof, and the solvates thereof and the hydrates thereof; and a use of a substance selected from the group consisting of the pyrimidone derivatives of formula (I) and the physiologically acceptable salts thereof, and the solvates thereof and the hydrates thereof for the manufacture of the aforementioned medicament.

[0011] As used herein, the C₁₋₆ alkyl group represents a straight or branched alkyl group having 1 to 6 carbon atoms, for example, methyl group, ethyl group, n-propyl group, isopropyl group, n-butyl group, isobutyl group, sec-butyl group, tert-butyl group, n-pentyl group, isopentyl group, neopentyl group, 1,1-dimethylpropyl group, n-hexyl group, isohexyl group, and the like;

The C₁₋₆ alkoxy group represents an alkyloxy group having from 1 to 6 carbon atoms, for example, methoxy group, ethoxy group, propoxy group, isopropoxy group, butoxy group, isobutoxy group, sec-butoxy group, tert-butoxy group, pentyloxy group, isopentyloxy group, neopentyloxy group, 1,1-dimethylpropyloxy group;

The halogen atom represents a fluorine, chlorine, bromine or iodine atom;

The C₁₋₂ perhalogenated alkyl group represents an alkyl group wherein all the hydrogen have been substituted by a halogeno, for example a CF₃ or C₂F₅,

The C₁₋₃ halogenated alkyl group represents an alkyl group wherein at least one hydrogen has not been substituted by a halogen atom.

[0012] The compounds represented by the aforementioned formula (I) may form a salt. Examples of the salt include, when an acidic group exists, salts of alkali metals and alkaline earth metals such as lithium, sodium, potassium, magnesium, and calcium; salts of ammonia and amines such as methylamine, dimethylamine, trimethylamine, dicyclohexylamine, tris(hydroxymethyl)aminomethane, N,N-bis(hydroxyethyl)piperazine, 2-amino-2-methyl-1-propanol, ethanolamine, N-methylglucamine, and L-glucamine; or salts with basic amino acids such as lysine, hydroxylysine, and arginine. The base-addition salts of acidic compounds are prepared by standard procedures well known in the art.

[0013] When a basic group exists, examples include salts with mineral acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid; salts with organic acids such as methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, acetic acid, propionic acid, tartaric acid, fumaric acid, maleic acid, malic acid, oxalic acid, succinic acid, citric acid, benzoic acid, mandelic acid, cinnamic acid, lactic acid, glycolic acid, glucuronic acid, ascorbic acid, nicotinic acid, and salicylic acid; or salts with acidic amino acids such as aspartic acid, and glutamic acid.

[0014] The acid-addition salts of the basic compounds are prepared by standard procedures well known in the art which include, but are not limited thereto, dissolving the free base in an aqueous alcohol solution containing the appropriate acid and isolating the salt by evaporating the solution, or by reacting the free base and an acid in an organic solvent, in which case the salt separates directly, or is precipitated with a second organic solvent, or can be obtained by concentration of the solution. The acids which can be used to prepare the acid-addition salts include preferably those which produce, when combined with the free base, pharmaceutically-acceptable salts, that is, salts whose anions are relatively innocuous to the animal organism in pharmaceutical doses of the salts, so that the beneficial properties inherent in the free base are not compromised by side effects ascribable to the anions. Although medicinally acceptable salts of the basic compounds are preferred, all acid-addition salts are within the scope of the present invention.

[0015] In addition to the pyrimidone derivatives represented by the aforementioned formula (I) and salts thereof, their solvates and hydrates also fall within the scope of the present invention. The pyrimidone derivatives represented by the aforementioned formula (I) may have one or more asymmetric carbon atoms. As for the stereochemistry of such asymmetric carbon atoms, they may independently be in either (R) and (S) configuration, and the pyrimidone derivative may exist as stereoisomers such as optical isomers, or diastereoisomers. Any stereoisomers in pure form, any mixtures of stereoisomers, racemates and the like fall within the scope of the present invention.

[0016] Furthermore, as the pyrimidone derivatives represented by the aforementioned formula (I), a 3H-4-one compound, a 4-hydroxy compound, and a 1H-4-one compound of may exist as tautomers. The existence of such tautomers is readily apparent to those skilled in the art, and these tautomers fall within the scope of the present invention.

[0017] Examples of particularly preferred compounds of the present invention are shown in table 1 thereafter and are listed hereabove. However, the scope of the present invention is not limited by these compounds.

[0018] Preferred compounds of the present invention represented by formula (I) include also :

(1) Compounds wherein R1 represents a 3- or 4-pyridyl group and more preferably 4-pyridyl group, which may be substituted by a C₁₋₂ alkyl group, C₁₋₂ alkoxy group or a halogen atom.

[0019] More preferred compounds of the present invention represented by formula (I) include also :

- (1) Compounds wherein R1 represents an unsubstituted 4-pyridyl group ;
- (2) Compounds wherein the heterocyclic ring is selected from :

2,3,4,9-tetrahydro-1H-pyrido[3,4-b]indole,
1,2,3,4-tetrahydroisoquinoline and
4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
1,2,3,4-tetrahydrobenzofuro[2,3-c]pyridine.

[0020] Particularly preferred compounds of the present invention represented by formula (I) include:

6-pyridin-4-yl-2-[2,3,4,9-tetrahydro-1H-pyrido[3,4-b]indol-2-yl]pyrimidin-4(1H)-one,
6-pyridin-4-yl-2-[1,2,3,4-tetrahydroisoquinolin-2-yl]pyrimidin-4(1H)-one,
6-pyridin-4-yl-2-[6,7-dimethoxy-1,2,3,4-tetrahydroisoquinolin-2-yl]pyrimidin-4(1H)-one,
6-pyridin-4-yl-2-[4,5,6,7-tetrahydrothieno[2,3-c]pyridin-6-yl]pyrimidin-4(1H)-one
and 6-pyridin-4-yl-2-[1,2,3,4-tetrahydrobenzofuro[2,3-c]pyridin-2-yl]pyrimidin-4(1H)-one.


[0021] As a further object, the present invention concerns also methods for preparing the pyrimidone compounds represented by the aforementioned formula (I).

[0022] These compounds can be prepared, for example, according to methods explained below.

Preparation method

[0023] Pyrimidone compounds represented by the aforementioned formula (I) may be prepared according to scheme 1.

Scheme 1




(In the above scheme the definition of R1 is the same as already described above).

[0024] The 2-methylthio derivative represented by the above formula (II), wherein R1 is as defined for compound of formula (I), is allowed to react with the heterocycle of formula (III) to obtain the compound of the aforementioned formula (I). The reaction may be carried out in a solvent such as, for example, an alcoholic solvent such as n-pentanol or isoamyl alcohol at a suitable temperature ranging from 100 - 180 °C under ordinary air.

[0025] Compound of formula (II) may be prepared according to the method defined in scheme 2.

Scheme 2



(In the above scheme R represents an alkyl group and the definition of R1 is the same as already described above).

[0026] According to this method, the 3-ketoester of formula (IV) is allowed to react with a 2-methyl-2-thiopseudourea sulfate. The reaction may be carried out in solvent such as water or an alcohol, such as ethanol, propanol and butanol, at a suitable temperature ranging from 25-100°C under ordinary air.

[0027] Compounds of formula (IV) and formula (II) are commercially available or may be synthesized according to known methods of one skilled in the art.

[0028] For example compounds of formula (II), wherein R1 represent a 4-pyridyl group optionally substituted by a C₁₋₄ alkyl group, C₁₋₄ alkoxy group or a halogen atom, can be prepared by reacting a nicotinic acid optionally substituted by a C₁₋₄ alkyl group, C₁₋₄ alkoxy group or an halogen, with a malonic acid monoester. The reaction can be carried out using methods well known to one skilled in the art, such as for example in presence of a coupling agent such as 1,1'-carbonylbis-1H-imidazole in a solvent such as a tetrahydrofuran at a temperature ranging from 20 to 70°C.

[0029] In addition when applicable, compound of formula (I) wherein the heterocyclic ring is substituted by a hydroxyl group, it can be alkylated to give a C₁₋₆ alkoxy group to obtain an other compound of formula (I), using well known methods in the art.

[0030] In the above reactions, protection or deprotection of a functional group may sometimes be necessary. A suitable protecting group can be chosen depending on the type of a functional group, and a method described in the literature may be applied. Examples of protecting groups, of protection and deprotection methods are given for example in *Protective groups in Organic Synthesis* Greene et al., 2nd Ed. (John Wiley & Sons, Inc., New York).

[0031] The compounds of the present invention have inhibitory activity against GSK3β. Accordingly, the compounds of the present invention are useful as an active ingredient for the preparation of a medicament, which enables preventive and/or therapeutic treatment of neurodegenerative diseases such as Alzheimer's disease. In addition, the compounds of the present invention are also useful as an active ingredient for the preparation of a medicament for preventive and/or therapeutic treatment of Parkinson's disease, frontoparietal dementia, corticobasal degeneration, Pick's disease, cerebrovascular accidents, brain and spinal cord trauma, and peripheral neuropathies.

[0032] The present invention further relates to a method for treating neurodegenerative diseases caused by abnormal activity of GSK3β and of the aforementioned diseases which comprises administering to a mammalian organism in need thereof an effective amount of a compound of the formula (I).

[0033] As the active ingredient of the medicament of the present invention, a substance may be used which is selected from the group consisting of the compound represented by the aforementioned formula (I) and pharmacologically acceptable salts thereof, and solvates thereof and hydrates thereof. The substance, per se, may be administered as the medicament of the present invention, however, it is desirable to administer the medicament in a form of a pharmaceutical composition which comprises the aforementioned substance as an active ingredient and one or more of pharmaceutical additives. As the active ingredient of the medicament of the present invention, two or more of the aforementioned substances may be used in combination. The above pharmaceutical composition may be supplemented

with an active ingredient of another medicament for the treatment of the above mentioned diseases. A type of the pharmaceutical composition is not particularly limited, and the composition may be provided as any formulation for oral or parenteral administration. For example, the pharmaceutical composition may be formulated, for example, in the form of pharmaceutical compositions for oral administration such as granules, fine granules, powders, hard capsules, soft capsules, syrups, emulsions, suspensions, solutions and the like, or in the form of pharmaceutical compositions for parenteral administrations such as injections for intravenous, intramuscular, or subcutaneous administration, drip infusions, transdermal preparations, transmucosal preparations, nasal drops, inhalants, suppositories and the like. Injections or drip infusions may be prepared as powdery preparations such as in the form of lyophilized preparations, and may be used by dissolving just before use in an appropriate aqueous medium such as physiological saline. Sustained-release preparations such as those coated with a polymer may be directly administered intracerebrally.

[0034] Types of pharmaceutical additives used for the manufacture of the pharmaceutical composition, content ratios of the pharmaceutical additives relative to the active ingredient, and methods for preparing the pharmaceutical composition may be appropriately chosen by those skilled in the art. Inorganic or organic substances, or solid or liquid substances may be used as pharmaceutical additives. Generally, the pharmaceutical additives may be incorporated in a ratio ranging from 1% by weight to 90% by weight based on the weight of an active ingredient.

[0035] Examples of excipients used for the preparation of solid pharmaceutical compositions include, for example, lactose, sucrose, starch, talc, cellulose, dextrin, kaolin, calcium carbonate and the like. For the preparation of liquid compositions for oral administration, a conventional inert diluent such as water or a vegetable oil may be used. The liquid composition may contain, in addition to the inert diluent, auxiliaries such as moistening agents, suspension aids, sweeteners, aromatics, colorants, and preservatives. The liquid composition may be filled in capsules made of an absorbable material such as gelatin. Examples of solvents or suspension mediums used for the preparation of compositions for parenteral administration, e.g. injections, suppositories, include water, propylene glycol, polyethylene glycol, benzyl alcohol, ethyl oleate, lecithin and the like. Examples of base materials used for suppositories include, for example, cacao butter, emulsified cacao butter, lauric lipid, witepsol.

[0036] Dose and frequency of administration of the medicament of the present invention are not particularly limited, and they may be appropriately chosen depending on conditions such as a purpose of preventive and/or therapeutic treatment, a type of a disease, the body weight or age of a patient, severity of a disease and the like. Generally, a daily dose for oral administration to an adult may be 0.01 to 1,000 mg (the weight of an active ingredient), and the dose may be administered once a day or several times a day as divided portions, or once in several days. When the medicament is used as an injection, administrations may preferably be performed continuously or intermittently in a daily dose of 0.001 to 100 mg (the weight of an active ingredient) to an adult.

Chemical Examples

[0037] The present invention will be explained more specifically with reference to the following general examples, however, the scope of the present invention is not limited to these examples.

Example 1 : Preparation of 2-("heterocycle")-6-pyridin-4-ylpyrimidin-4(1H)-one.

1.1. Preparation of Ethyl 3-(4-pyridyl)-3-oxopropionate

[0038] Isonicotinic acid (35.56 g, 289 mmol) was added to a solution of 1,1'-carbonylbis-1H-imidazole (46.98 g, 290 mmol) in tetrahydrofuran (700ml), and the resulting solution was stirred for 1.5 hr at 50°C. After cooling to room temperature, malonic acid monoester potassium salt (51.7 g, 304 mmol) and magnesium chloride (34.33 g, 361 mmol) were added, and the mixture was refluxed for 1 hr and then heated at 50°C for 6 hr. The solvent was removed under reduced pressure and the residue was quenched by the addition of dilute acetic acid. The organic layer was extracted with ethyl acetate (3 times) and the combined extracts were washed with dilute aqueous sodium bicarbonate and brine, and were concentrated under reduced pressure. The residue was purified by silica gel column chromatography (eluent; hexane/ethyl acetate = 2/1 to 1/1) and recrystallization from hexane - ethyl acetate gave 41.52 g (74%) of the title compound.

1.2 Preparation of 2-(Methylthio)-6-pyridinyl-4-ylpyrimidin-4(1 H)-one

[0039] To a solution of 5.76 g (20.7 mmol) of 2-methyl-2-thiopseudoiurea sulfate in 48 ml of water was added 4.85 g (86.52 mmol) of potassium hydroxide. The mixture was agitated and 8.0 g (41.4 mmol) of ethyl 3-(4-pyridyl)-3-oxopropionate was added and stirring was maintained for 48 hours. The precipitate was recovered by filtration and was washed with water and then ether. The product was dried at 90°C in vacuo to give 6.26 g, 69% of white solid.


Mp : 328-330°C.

1.3 Preparation of 2-["heterocycle"]-6-pyridin-4-ylpyrimidin-4(1H)-one

[0040] A solution of 1 equivalent of 2-(methylthio)-6-pyridinyl-4-ylpyrimidin-4(1H)-one and 3 to 5 equivalents of a (hetero)aryl-tetrahydro pyridine of formula (III) were suspended in amyl alcohol (0.1 - 0.2 M) and was heated at 150°C during 72 hours. The cooled solution was concentrated and purified by chromatography on silica gel.

[0041] A list of chemical structures and physical data for compounds of the aforementioned formula (I) illustrating the present invention is given in Table 1.

Table 1



(I)

R1 = 4-pyridyl

No.	Structure	m.p.(°C)	[M+H] ⁺
1		-	311
2		-	344
3		-	305
4		240.1-240.5	365
5		-	345

EP 1 136 493 A1

Test Example: Inhibitory activity of the medicament of the present invention against GSK3 β :

[0042] Two different protocols can be used.

[0043] In a first protocol : 7.5 μ M of prephosphorylated GS1 peptide and 10 μ M ATP (containing 300,000 cpm of 33P-ATP) were incubated in 25 mM Tris-HCl, pH 7.5, 0.6 mM DTT, 6 mM MgCl₂, 0.6 mM EGTA, 0.05 mg/ml BSA buffer for 1 hour at room temperature in the presence of GSK3 β (total reaction volume : 100 microliters).

[0044] In a second protocol : 4.1 μ M of prephosphorylated GS1 peptide and 42 μ M ATP (containing 260,000 cpm 33P-ATP) were incubated in 80 mM Mes-NaOH, pH 6.5, 1 mM Mg acetate, 0.5 mM EGTA, 5 mM 2-mercaptoethanol, 0.02% Tween 20, 10% glycerol buffer for 2 hours at room temperature in the presence of GSK3 β . Inhibitors were solubilised in DMSO (final solvent concentration in the reaction medium, 1%).

[0045] The reaction was stopped with 100 microliters of a solution made of 25 g polyphosphoric acid (85% P₂O₅), 126 ml 85% H₃PO₄, H₂O to 500 ml and then diluted to 1 :100 before use. An aliquot of the reaction mixture was then transferred to Whatman P81 cation exchange filters and rinsed with the solution described above. Incorporated 33P radioactivity was determined by liquid scintillation spectrometry.

The phosphorylated GS-1 peptide had the following sequence :



[0046] The GSK3 β inhibitory activity of the compounds of the present invention are expressed in IC₅₀, and as an illustration the range of IC₅₀'s of the compounds given in table 1, is between 0.3 to 10 micromolar concentrations.

Formulation Example

(1) Tablets

[0047] The ingredients below were mixed by an ordinary method and compressed by using a conventional apparatus.

Compound of Example 1	30 mg
Crystalline cellulose	60 mg
Corn starch	100 mg
Lactose	200 mg
Magnesium stearate	4 mg

(2) Soft capsules

[0048] The ingredients below were mixed by an ordinary method and filled in soft capsules.

Compound of Example 1	30 mg
Olive oil	300 mg
Lecithin	20 mg

(3) Parenteral preparations

[0049] The ingredients below were mixed by an ordinary method to prepare injections contained in a 1 ml ampoule.


Compound of Example 1	3 mg
Sodium chloride	4 mg
Distilled water for injection	1 ml

Industrial Applicability

[0050] The compounds of the present invention have GSK3 β inhibitory activity and are useful as an active ingredient of a medicament for preventive and/or therapeutic treatment of neurodegenerative diseases caused by abnormal activity of GSK3 β .


Claims

1. A pyrimidone derivative represented by formula (I) or a salt thereof, or a solvate thereof or a hydrate thereof:




wherein:

- 20
- R1 represents a pyridyl ring optionally substituted by a C₁₋₄ alkyl group, C₁₋₄ alkoxy group or a halogen atom ;
 - The heterocyclic ring of formula :



is selected from group consisting of:



the heterocyclic ring being optionally substituted by 1 to 3 substituents selected from a C₁₋₆ alkyl group, halogeno, a C₁₋₂ perhalogenated alkyl group, a C₁₋₃ halogenated alkyl group, a hydroxyl group; a C₁₋₆ alkoxy group, methylenedioxy group, a nitro, a cyano, or a phenyl ring; and R2 represents a hydrogen atom or a C₁₋₆ alkyl group.

- 50
2. A pyrimidone derivative or a salt thereof, or a solvate thereof or a hydrate thereof according to claim 1, wherein R1 represents an unsubstituted 4-pyridyl group.
- 55
3. A pyrimidone derivative or a salt thereof, or a solvate thereof or a hydrate thereof according to claim 2, wherein the heterocyclic ring is selected from:

2,3,4,9-tetrahydro-1H-pyrido[3,4-b]indole,
1,2,3,4-tetrahydroisoquinoline and

EP 1 136 493 A1

4,5,6,7-tetrahydrothieno[2,3-*c*]pyridine.
4,5,6,7-tetrahydrofuro[2,3-*c*]pyridine.
1,2,3,4-tetrahydrobenzofuro[2,3-*c*]pyridine.

- 5 4. A pyrimidone derivative which is selected from the group consisting of:

6-pyridin-4-yl-2-[2,3,4,9-tetrahydro-1*H*-pyrido[3,4-*b*]indol-2-yl]pyrimidin-4(1*H*)-one,
6-pyridin-4-yl-2-[1,2,3,4-tetrahydroisoquinolin-2-yl]pyrimidin-4(1*H*)-one,
6-pyridin-4-yl-2-[6,7-dimethoxy-1,2,3,4-tetrahydroisoquinolin-2-yl]pyrimidin-4(1*H*)-one,
10 6-pyridin-4-yl-2-[4,5,6,7-tetrahydrothieno[2,3-*c*]pyridin-6-yl]pyrimidin-4(1*H*)-one, and
6-pyridin-4-yl-2-[1,2,3,4-tetrahydrobenzofuro[2,3-*c*]pyridin-2-yl]pyrimidin-4(1*H*)-one,
or a salt thereof, or a solvate thereof or a hydrate thereof.

- 15 5. A medicament comprising as an active ingredient a substance selected from the group consisting of a pyrimidone derivative according to claim 1.

6. A GSK3 β inhibitor selected from the group of a pyrimidone derivative according to claim 1.

- 20 7. Use of a compound according to claim 1 for the preparation of a medicament for preventive and/or therapeutic treatment of a disease caused by abnormal GSK3 β activity.

8. Use of a compound according to claims 1 to 4 for the preparation of a medicament for preventive and/or therapeutic treatment of a neurodegenerative disease.

- 25 9. Use of a compound according to claim 8, wherein the disease is selected from the group consisting of Alzheimer's disease, Parkinson's disease, frontoparietal dementia, corticobasal degeneration, Pick's disease, cerebrovascular accidents, brain and spinal cord trauma, and peripheral neuropathies.